



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Driftwood from the arctic sea.—A large amount of driftwood has been examined by INGVARSON,²³ from Spitzbergen, Beeren Island, Jan Mayen, Greenland, and the mouth of Jenisei River. Among the conifers are species of *Larix*, *Picea*, *Pinus*, and *Abies*; while the deciduous plants belong mostly to *Salix*, *Populus*, and *Betula*. The structure of the trunks and roots is described and figured. The author reaches the conclusion that the polar current is a much more important factor in the transportation of driftwood in the arctic seas than is the gulf stream. The fact that seeds and fruits of various arctic plants have been found caught in the fissures of driftwood leads to the conclusion that the wide distribution of several arctic species may be due to such transportation.—THEO. HOLM.

A mutant of *Leptospermum scoparium*.—This shrub is very common in New Zealand, occupying vast tracts of usually barren country. The flowers are white, and the close-growing shrubs when in full bloom look as if densely powdered with snow. Pink forms are found now and then, but COCKAYNE²⁴ records the sudden appearance of a single plant with rich crimson flowers. This plant was taken possession of by a firm of nurserymen, and among the progeny obtained from the seed six more plants with crimson flowers appeared. The variation extends also to the size and color of the leaves, which are brownish instead of green.—J. M. C.

***Linnaea borealis*.**—WITTRÖCK²⁵ has published thirteen mostly colored plates of Swedish forms of this species, and in the accompanying text has described about 160 forms. The vegetative organs are mentioned, but the author has made a special study of the shape and coloration of the corolla, which is beautifully illustrated. The American representatives are briefly described, and the following may be noted: *L. borealis longiflora* Torr. f. *angustissima* (Washington), f. *curticalyx* (Colorado), f. *minutifolia* (Vermont), f. *integerrima* (Alberta).—THEO. HOLM.

Monograph of *Fraxinus*.—LINGELSHEIM²⁶ has made a general study of *Fraxinus* as to its morphology (in the old sense), anatomy, geographical distribution, fossil forms, phylogeny, and classification. He recognizes twenty-seven species in § *Ornus*, nine of which are new; and thirty-two species in § *Fraxinaster*, four of which are new. The four new *Fraxinaster*s are *F. rufescens* (Jarl 1315), *F. papillosa* (Townsend & Barber 354), *F. Pringlei* (Pringle 9417), all from Mexico, and *F. hybrida* from Florida (Curtiss 2321, Nash 941 and 1698).—J. M. C.

²³ INGVARSON, FR., Om Drifverden i Norra Ishafvet. K. Vet. Akad. Hdlgr. 37:1. 1903.

²⁴ COCKAYNE, L., On the sudden appearance of a new character in an individual of *Leptospermum scoparium*. New Phytol. 6:43-46. 1907.

²⁵ WITTRÖCK, V. BR., *Linnaea borealis* L., species polymorpha et polychroma. Acta. Hort. 4:no. 7. 1907.

²⁶ LINGELSHEIM, ALEXANDER, Vorarbeiten zu einer Monographie der Gattung *Fraxinus*. Bot. Jahrb. 46:185-223. pl. 8. 1907.